

ON ORDER CONTINUOUS BANACH $C(K)$ -MODULES

Omer Gok

Yildiz Technical University
Faculty of Arts and Sciences
Mathematics Department, Davutpasa
Istanbul - 34210, TURKEY

Abstract: Let X be a Banach space and let X' be the norm dual of X . $C(K)$ denotes the Banach space of all continuous real or complex valued functions on a compact Hausdorff space K with the supremum norm. Suppose that X is a Banach $C(K)$ -module. In this paper, we characterize the order continuity of Banach $C(K)$ -module if dual Banach space X' does not contain the space ℓ^∞ of all bounded sequences by using Arens multiplication.

AMS Subject Classification: 46H25, 46B42, 47B60

Key Words: Banach lattices; Arens multiplication; order continuous norm; cyclic vector; center of Banach lattices

1. Introduction

Lozanovsky's study in [9] is that a Dedekind complete Banach lattice has order continuous norm if and only if it does not contain a copy of ℓ^∞ . Kitover and Orhon [7] investigate that firstly what is an analogue of Dedekind completeness for Banach $C(K)$ -modules and secondly, what should be considered as an analogue of order continuity for Banach $C(K)$ -modules, [11]. The positive answer to the question in the first case is provided by the well-known notion of Kaplansky module by Kitover and Orhon [7]. Positive answer to the second assertion is given again by Kitover and Orhon [7].

In this paper, we present the dual versions of these solutions. Namely, we take dual Banach $C(K)$ -modules instead of Banach $C(K)$ -modules.

For unexplained notion and terminology in this paper, we refer to the books [1] and [8], [10].

Definition 1. ([1]) Let X be a Banach space and let $C(K)$ be the Banach space of all real or complex valued continuous functions on a compact Hausdorff space K . A bilinear mapping $b : C(K) \times X \rightarrow X$ is called a Banach $C(K)$ -module if the following conditions are satisfied:

- i.a. $(b.x) = (a.b).x$ for all $a, b \in C(K)$ and $x \in X$,
- ii. $1.x = x$ for all $x \in X$, where 1 is the constant one function,
- iii. The mapping b is continuous, i.e., the inequality

$$\|a.x\| \leq \|a\|\|x\|$$

holds for every $a \in C(K)$ and $x \in X$.

We can extend the bilinear mapping b to dual spaces X' and X'' by using the Arens multiplication, [2] and [3]:

$$b' : X' \times X \rightarrow C(K)'$$

is defined by $b'(x', x)(a) = x'(b(a, x))$.

$$b'' : C(K)'' \times X' \rightarrow X'$$

is defined by $b''(\hat{a}, x')(x) = \hat{a}(b'(x', x))$.

$$b''' : C(K)'' \times X'' \rightarrow X''$$

is defined by $b'''(\hat{a}, x'')(x') = x''(b''(\hat{a}, x'))$.

In these mappings, if we put $X = C(K)$, then $C(K)'' = C(S)$ where S is a hyperstonian compact Hausdorff space. Let B be the Boolean algebra of all idempotents in $C(S)$. By bilinear mapping b defines a continuous map $m : C(K) \rightarrow L(X)$ defined by $m(a)x = a.x = b(a, x)$. Also, bilinear mapping b'' defines a continuous mapping $m^* : C(K)'' \rightarrow L(X')$ defined by $m^*(a)x' = a.x'$.

The properties of the mappings m and m^* are given by the following theorem.

Theorem 2. ([5]) *Let X be a Banach $C(K)$ -module and let $m : C(K) \rightarrow L(X)$ be a continuous mapping from $C(K)$ to the space $L(X)$ of all bounded linear operators from X into X . Then, the following claims are true:*

- i. m is an algebra homomorphism.
- ii. The mapping m^* satisfies: $m^*(a) = (m(a))^*$ for all $a \in C(K)$, where $(m(a))^*$ is the adjoint of $m(a)$.
- iii. $m(1) = I$, where I denotes the identity operator.
- iv. $m^* : C(K)'' \rightarrow L(X')$ is a continuous mapping defined by $m^*(a)(x') = a.x'$.
- v. X' is a Banach $C(K)''$ -module.
- vi. X'' is a Banach $C(K)''$ -module.
- vii. m^* is a weak* to weak* continuous mapping from $C(K)''$ to X' .
- viii. m^* is an algebra homomorphism.

Proof. Proofs are done directly by using the Arens extension mappings and module. □

Let X be a Banach $C(K)$ -module. Then, we can define an equivalent norm on X' ,

$$\|x'\|_1 = \sup\{\|m^*(a)x'\| : a \in C(S), \|a\| \leq 1\}.$$

The homomorphism m^* is a contraction with respect to new norm. The $kerm^*$ is a closed ideal in $C(S)$, because of algebra homomorphism m^* . We can replace $C(S)$ by $C(T) \cong C(S)/kerm^*$, where $T = \{t \in S : a(t) = 0 \forall a \in kerm^*\}$. So, we can assume that m^* is one to one, [4].

Theorem 3. ([4, 5]) *Let X be a Banach $C(K)$ -module and let $m^* : C(K)'' \rightarrow L(X')$ be a contractive homomorphism. Then,*

- i. *If for every $a, b \in C(K)''$, $|a| \leq |b|$, then $\|m^*(a)x'\| \leq \|m^*(b)x'\|$ for any $x' \in X'$.*
- ii. *If m^* is one to one, then it is an isometry.*

Definition 4. ([5, 7]) *Let $m : C(K) \rightarrow L(X)$ be a bounded unital algebra homomorphism and let $x \in X$. An idempotent $e_x \in B$ is said to be a carrier projection of x if $m(e_x)x = x$ and $e_x \leq e$ in $C(K)$, whenever $e \in B, m(e)x = x$.*

Theorem 5. ([7]) *Let X be a Banach $C(K)$ -module and let $m^* : C(K)'' \rightarrow L(X')$ be a continuous unital algebra homomorphism. Let $x' \in X'$ and $e_{x'}$ be the carrier projection of x' . Then, for any $a \in C(K)''$ and $m^*(a)x' = 0$ if and only if $ae_{x'} = 0$.*

Proof. Let $ae_{x'} = 0$. Then,

$$m^*(a)x' = m^*(a)m^*(e_{x'})x' = m^*(ae_{x'})x' = 0.$$

Assume that for some $e \in B, m^*(e)x' = 0$. Then, $m^*((1 - e)x') = x'$ and therefore $e_{x'} \leq 1 - e$. So, $ee_{x'} = 0$. Suppose that for some $0 \leq a \in C(K)''$. We have $m^*(a)x' = 0$ and that for some $t \in S, a(t) > 0$. Since S is totally disconnected, there are some $\epsilon > 0$ and $e \in B$ such that $e(t) = 1$ and $0 \leq \epsilon e \leq a$. From here we get $\epsilon \|m^*(e)x'\| \leq \|m^*(a)x'\| = 0$. Therefore, $ee_{x'} = 0$ and $e_{x'}(t) = 0$ and hence $ae = 0$.

If $m^*(a)x' = 0$ for some $a \in C(K)''$, then $m^*(|a|)x' = 0$ hence $|a|e_{x'} = 0$ and so $ae_{x'} = 0$. □

Definition 6. ([7]) A Banach space $C(K)$ -module X is called a Veksler module if any $x \in X \setminus \{0\}$ has a carrier projection $e_x \in C(K)$.

A compact Hausdorff space K is called quasi-Stonian if it is basically disconnected, that is, the closure of every open G_δ set in K is open. It is well-known that the following claims are equivalent:

- a. K is quasi-Stonian,
- b. $C(K)$ is σ -Dedekind complete,
- c. Every non-negative sequence bounded from above in $C(K)$ has a supremum in $C(K)$.
- d. Every principal band in $C(K)$ is a projection band.

Definition 7. ([7]) A Banach $C(K)$ -module X is said to be a Kaplansky module if it satisfies the following conditions:

- i. The compact space K is Stonian,
- ii. For any $x \in X$ and for any non-negative set $\{a_\alpha\}$ bounded above in $C(K)$ the following holds: if $a_\alpha x = 0$ for all α , then $ax = 0$, where $a = \sup_\alpha a_\alpha$.

Theorem 8. ([6, 7, 9]) *Let E be a σ -Dedekind complete Banach lattice. The following are equivalent:*

- i. *The original lattice norm on E is order continuous.*
- ii. *E does not contain ℓ^∞ as a closed subspace.*
- iii. *E does not contain ℓ^∞ as a closed sublattice.*

Definition 9. ([5, 6, 7]) Let B be a Boolean algebra of projections in $L(X)$. B is called a Bade-complete Boolean algebra of projections if for any

increasing net (e_α) in B and for every $x \in X$ we have $\lim_\alpha \|(e - e_\alpha)x\| = 0$, where $e = \sup_\alpha e_\alpha$.

Let B be idempotents in $C(K)'' = C(S)$. B consists of characteristic functions of the clopen subsets of S . And $m^*(B) = B^*$ is a Bade -complete Boolean algebra of projections on X' .

2. The cyclic Banach spaces

In this section we define cyclic Banach space by using Banach $C(K)$ -module.

Definition 10. ([1, 7]) Let X be a Banach $C(K)$ -module and $x \in X$. The cyclic subspace $X(x)$ of X is defined by

$$X(x) = Cl\{m(a)x : a \in C(K)\},$$

where the notation Cl denotes the closure of a set.

A Banach $C(K)$ -module X is said to be a cyclic Banach space if there is an $x \in X$ such that $X = X(x)$.

Similar definition is given on dual Banach space X' .

Definition 11. The cyclic subspace $X'(x')$ of X' is defined by

$$X'(x') = Cl\{m^*(a)x' : a \in C(S)\},$$

where the Cl denotes the closure.

The Banach $C(S)$ -module X' is called a cyclic Banach space if there exists a vector $x' \in X'$ such that $X' = X'(x')$.

Let X be a Banach $C(K)$ -module. Then, it defines a unital bounded homomorphism $m : C(K) \rightarrow L(X)$. By means of the Arens product, we define a unital bounded homomorphism $m^* : C(K)'' \rightarrow L(X')$. Assume $X' = X'(x'_0)$ for some vector $x'_0 \in X'$, that is, X' is a cyclic Banach space and x'_0 is a cyclic vector. Then, the following assertions are true, [6]:

- i. X' can be represented as a Banach lattice with quasi-interior point x'_0 .
- ii. The cone of X' is identified by the set

$$(X')^+ = Cl\{m^*(a)x' : 0 \leq a \in C(S)\}.$$

iii. The center $Z(X')$ of the Banach lattice X' is the weak* closure of $m^*(C(S))$.

iv. The unit ball of $Z(X')$ is the closure of unit ball of $m^*(S)$ in the weak* operator topology.

v. If x' is quasi-interior point in the Banach lattice X' , then for the order ideal $A_{x'}$ generated by x' we have $A_{x'} = Z(X')x'$.

We give the following theorem concerning cyclic Banach spaces.

Theorem 12. *Assume that X' is a cyclic Banach $C(S)$ -module, x'_0 is a cyclic vector in X' and let B be the Boolean algebra of all idempotents in $C(S)$. Then,*

$m^(C(S))$ is weak* operator closed and X' has an order continuous norm if it is represented as a Banach lattice.*

Proof. Since the compact space S is Stonian and B is a Bade complete Boolean algebra of projections on X' , the conclusion is true. \square

Theorem 13. ([7]) *Let X be a Banach $C(K)$ -module and let $m^* : C(K)'' \rightarrow L(X')$ be an injective bounded unital algebra homomorphism such that $m^*(C(S))$ is a weak*operator closed in $L(X')$. The following assertions are equivalent:*

i. *X' is a Veksler module and no cyclic subspace of X' contains a copy of l^∞ .*

ii. *Each cyclic subspace of X' has order continuous norm when it is represented as a Banach lattice.*

iii. *X' is a Kaplansky module and no cyclic subspace of X' contains a copy of l^∞ .*

Proof. *iii* \Rightarrow *i*. Every Kaplansky module is a Veksler module.

i \Rightarrow *iii*. If X' is represented as a Banach lattice with a quasi-interior point x'_0 , then it is σ -Dedekind complete and $m^*(C(S)) = Z(X')$. So, $m^*(C(S))$ is weak* operator closed. Since X' does not contain any copy of l^∞ , X' has order continuous norm.

iii \Rightarrow *ii*. Since X' has order continuous norm, X' is Dedekind complete. Hence, it does not contain a copy of l^∞ . \square

References

- [1] Yu.A. Abramovich, E.L. Arenson, A.K. Kitover, *Banach $C(K)$ -Modules and Operators Preserving Disjointness*, Pitman Research Notes in Math. Ser., **277**, Longman Scientific and Technical (1992).
- [2] R. Arens, The adjoint of bilinear operations, *Proc. Amer. Math. Soc.*, **2** (1951), 839-848.
- [3] O. Gok, b-bimorphisms, *International Journal of Applied Mathematics*, **32**, No 1 (2019), 45-50; doi: 10.12732/ijam.v32i1.4.
- [4] D. Hadwin, M. Orhon, A noncommutative theory of Bade functionals, *Glasgow Math. J.*, **33** (1991), 75-81.
- [5] D. Hadwin, M. Orhon, Reflexivity and approximate reflexivity for Boolean algebras of projections, *J. Funct. Anal.*, **87** (1989), 348-353.
- [6] A. Kitover, M. Orhon, Reflexivity of Banach $C(K)$ -modules via reflexivity of Banach lattices, *Positivity*, **18**, No 3 (2014), 475-488.
- [7] A. Kitover, M. Orhon, Dedekind complete and order continuous Banach $C(K)$ -modules, In: *Positivity and Noncommutative Analysis*, Trends Math. Birkhauser-Springer (2019), 281-294.
- [8] P. Meyer-Nieberg, *Banach Lattices*, Springer, Berlin (1991).
- [9] G.Ya. Lozanovskii, On isomorphic Banach structure, *Siberian Math. J.*, **10**, No 1 (1969), 64-68.
- [10] H.H. Schaefer, *Banach Lattices and Positive Operators*, Springer, Berlin (1974).
- [11] W. Wnuk, *Banach Lattices with Order Continuous Norm*, Polish Scientific Publishers PWN, Warszawa (1999).

